

# Pain Management Card

If you ever need emergency medical treatment, your Pain Management Card lets other doctors and nurses know that you are taking VIVITROL®. This card includes information that will help them provide you with the best possible care.

If you are being treated with VIVITROL, it is important that you print this page, cut out the card below, fold in half and tape shut. Keep card with you at all times.



Print, cut out, fold in half and tape shut.

**Important Information For Emergency Pain Management**

I am currently taking VIVITROL® (naltrexone for extended-release injectable suspension), an opioid antagonist. Please see the back of this card for important information about pain management.

**My name:** \_\_\_\_\_

**Emergency contact name:** \_\_\_\_\_

**Emergency contact number:** \_\_\_\_\_

**My doctor:** \_\_\_\_\_

**My doctor's phone number:** \_\_\_\_\_

©2011 Alkermes, Inc. All rights reserved. VIV 701C January 2011 Printed in U.S.A.

**Call 1-888-235-8008**

For Full Prescribing Information, Including Boxed Warning, Please Visit [www.ntx-xrmedinfo.com](http://www.ntx-xrmedinfo.com)

**Vivitrol**  
(naltrexone for extended-release injectable suspension) 380 mg/vial

For Full Prescribing Information, Including Boxed Warning, Please Visit [www.ntx-xrmedinfo.com](http://www.ntx-xrmedinfo.com)

**Call 1-888-235-8008**

**To Medical Personnel Treating Me In An Emergency**

In an emergency situation in patients receiving VIVITROL, suggestions for pain management include regional analgesia or use of non-opioid analgesics. If opioid therapy is required as part of anesthesia or analgesia, patients should be continuously monitored in an anesthesia care setting by persons not involved in the conduct of the surgical or diagnostic procedure. The opioid therapy must be provided by individuals specifically trained in the use of anesthetic drugs and the management of the respiratory effects of potent opioids, specifically the establishment and maintenance of a patent airway and assisted ventilation. Irrespective of the drug chosen to reverse VIVITROL blockade, the patient should be monitored closely by appropriately trained personnel in a setting equipped and staffed for cardiopulmonary resuscitation.